

OPIOIDS: Prescribing, Withdrawal, and Medication Assisted Treatment

Program Test

In order to receive credit, please answer the entire program test, post evaluation form, and return via email to Gwen.Hunt@rwjbh.org or fax to 732-557-8935. Mark answers clearly.

1. Naloxone reverses only exogenous opioids: ___ True ___ False
2. 1 in how many people are expected to develop a substance abuse disorder in the U.S. over their lifetime?
 - a. 1 in 10
 - b. 1 in 20
 - c. 1 in 4
 - d. 1 in 7
3. Tolerance for the following symptoms never develop in chronic opioid users:
 - a. Respiratory Depression
 - b. Euphoria
 - c. Itching
 - d. Constipation
4. Buprenorphine combination therapy are sublingual films containing Buprenorphine and.....
 - a. Oxycodone
 - b. Naloxone
 - c. Naltroxone
 - d. Heroin
5. Buprenorphine can be started on a patient.....
 - a. While fully intoxicated with opioid in the ER
 - b. While fully intoxicated with alcohol in the ER
 - c. While in mild withdrawal from opioid in any clinical setting
 - d. While unresponsive due to opioid overdose

6. The risk of death associated with acute withdrawal of this substance is 10-25%
- Opioids
 - Marijuana
 - Alcohol
 - Cocaine
7. The leading cause of deaths in the U.S. currently is substance dependence on
- Opioids
 - Alcohol
 - Cocaine
 - Marijuana
8. Buprenorphine is safe in pregnancy: ___ True ___ False
9. Which of the following is true regarding Neonatal Abstinence Syndrome?
- It is self-limiting after birth, baby can go home with no care needed
 - It is life threatening to the baby, needs NICU monitoring
 - Baby can go home on OP narcotics
 - Can be avoided if the mother breast feeds the baby while continuing to use opioids
10. Buprenorphine is a schedule _____ drug
- I
 - II
 - III