

The Comprehensive Stroke Center: 2020 Performance and Quality Measures

Robert Wood Johnson University Hospital (RWJUH) Stroke Center Performance & Quality Measures

Clinical Measure	RWJ Goal	NJ Hospitals	RWJUH-NB
Early Antithrombotics: Patients with ischemic stroke or TIA who receive antithrombotic therapy by the end of hospital day two.	> 85%	97.7%	93.8%
VTE Prophylaxis: Patients with an ischemic stroke, or a hemorrhagic stroke, or stroke not otherwise specified who receive VTE prophylaxis the day of or the day after hospital admission.	> 85%	95%	99%
Antithrombotic: Percent of patients with an ischemic stroke or TIA prescribed antithrombotic therapy at discharge.	>85%	97.6%	99.6%
Statin: Ischemic stroke patients who are prescribed statin medication at hospital discharge	>85%	98.7%	99.1%
Anticoag for AFib/Aflutter: Percent of patients with an ischemic stroke or TIA with atrial fibrillation/flutter discharged on anticoagulation therapy.	>85%	97.7%	98.8%
Serious Complication Rates for Asymptomatic Carotid Endarterectomy (CEA) and Carotid Arterial Stenting (CAS)	<6%		0%
Time to IV Thrombolytic (tPA) Therapy within 45 min: Percent of acute ischemic stroke patients receiving IV tPA (clot busting medication) during the hospital stay who have a time from hospital arrival to initiation of tPA (door-to-needle) of 45 min or less.	>50%	60.1%	75.4%
Time of Door to Start of Device (DTD): DTD within 60 minutes for patients transferred from an outside hospital OR within 90 minutes for patients presenting directly (6 hour treatment Window)	>50%	29.3%	63.3%

This Get With The Guidelines® Aggregate Data report was generated using the Quintiles PMT® system. Copy or distribution of the Get With The Guidelines® Aggregate Data is prohibited without the prior written consent of the American Heart Association and Quintiles.